



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2009

RAS meets SLE

Wüthrich, R P

DOI: <https://doi.org/10.1093/ndt/gfp326>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-26516>

Journal Article

Published Version

Originally published at:

Wüthrich, R P (2009). RAS meets SLE. Nephrology, Dialysis, Transplantation, 24(9):2634-2636.

DOI: <https://doi.org/10.1093/ndt/gfp326>

RAS meets SLE*

Rudolf P. Wüthrich

Division of Nephrology, University Hospital, Zürich, Switzerland

Correspondence and offprint requests to: Rudolf P. Wüthrich; E-mail: rudolf.wuethrich@usz.ch

*Comment on Crowley SD, Vasievich MP, Ruiz P *et al.* Glomerular type 1 angiotensin receptors augment kidney injury and inflammation in murine autoimmune nephritis. *J Clin Invest* 2009; 119: 943–953

Keywords: renin-angiotensin system (RAS); lupus; inflammation

Role of the type 1 angiotensin receptor AT₁ in immune-mediated glomerular injury

The renin-angiotensin system (RAS) plays an important role in the pathogenesis of glomerular diseases. To delineate more precisely the role of RAS in creating renal injury, knockout mouse strains lacking the angiotensin type 1 or type 2 receptors (AT₁ and AT₂) have proven very informative.

Hence, in their elegant study that was published in the April 2009 issue of the *Journal of Clinical Investigation*, Crowley *et al.* have tested the effect of a targeted disruption of the AT₁ receptor on the glomerular disease process in a mouse lupus nephritis model [1]. MRL-*Fas^{lpr/lpr}* mice develop a severe lupus-like autoimmune syndrome that is characterized by severe proliferative immune-complex glomerulonephritis, autoantibody production and arthritis, in addition to a peculiar lymphoproliferative phenotype.

In their study, a unique MRL-*Fas^{lpr/lpr}* mouse strain that lacks the major type 1 angiotensin receptor (AT_{1A}) was generated to examine the role of angiotensin in the pathogenesis of autoimmune glomerular injury. Despite the absence of hypertension, AT_{1A} deficiency did not protect against disease but worsened kidney pathology and proteinuria, and accelerated mortality in these mice. Transplantation of AT_{1A}-deficient bone marrow into MRL-*Fas^{lpr/lpr}* did not affect survival, suggesting that the enhanced disease severity was not a consequence of immune cell alteration. Furthermore, AT_{1A}-deficient MRL-*Fas^{lpr/lpr}* mice did not display increased extrarenal tissue injury, and the lymphoproliferative phenotype as well as the circulating antinuclear antibodies were not changed.

How could this worsened disease in AT_{1A}-deficient MRL-*Fas^{lpr/lpr}* mice be explained? Unlike humans, rodents are known to express a second type 1 angiotensin recep-

tor (termed AT_{1B}), in particular in glomerular podocytes. Mice with AT_{1A} deficiency display a higher renin production and an enhanced activity of RAS. Evidence is then provided in this study for exaggerated AT_{1B} receptor activation in podocytes, manifesting as podocyte injury and increased production of inflammatory mediators. Treatment with the angiotensin receptor blocker (ARB) losartan that blocks both type 1 angiotensin receptors indeed reduced glomerular injury. Thus, it appears that the glomerular type 1 angiotensin receptor AT_{1B} mediated the augmented kidney injury and inflammation in AT_{1A}-deficient MRL-*Fas^{lpr/lpr}* mice. Altogether, the data suggest that an activation of AT₁ receptors may be sufficient to augment autoimmune glomerular injury and inflammation in the absence of hypertension.

SLE, hypertension and the RAS

In addition to the more direct effects of angiotensin II on the cardiovascular system, the RAS promotes immune and inflammatory tissue injury (Figure 1). Thus, it has been shown for example that activation of the type 1 angiotensin receptor AT₁ augments alloreactivity in experimental allograft transplantation [2]. Most of the actions of angiotensin II to enhance inflammation and tissue injury appear to be mediated by the AT₁ receptor. Activation of the opposing angiotensin receptor AT₂ is generally felt to mediate protection from inflammation [3]; however, in other situations, activation of AT₂ has also been linked to inflammation, such as increased chemokine expression in the kidney [4].

It is well known that the extent of hypertension has a major impact on target organ damage in human SLE [5]. Association studies in patients with lupus nephritis have linked high blood pressure to accelerated progression of renal failure. Case series have shown a general benefit of RAS inhibition in patients with SLE [6], but so far no controlled trials with ACEi or ARB have been performed. Although there are no such randomized trials in humans




Fig. 1. Pathophysiological alterations that are mediated by the different AT receptors. In humans, there is an AT₁ and an AT₂ receptor that have antagonizing functions. In rodents, there is an alternative AT₁ receptor termed AT_{1B} that is expressed in glomeruli by podocytes, and which is felt to have similar functions as the AT_{1A} receptor.

with SLE, it is felt that RAS inhibition confers protection in human lupus nephritis.

Animal models that apply pharmacological blockade of the RAS or targeted deletion of the angiotensin receptor genes have proved quite useful to gain a better understanding of how RAS promotes immune/inflammatory renal injury. Studies in lupus mice have revealed that inhibition of RAS with ACEi or ARB is beneficial, reducing glomerular injury and albuminuria, and TGF- β production, the latter being a major mediator of fibrosis [7,8]. However, the role of RAS in regulating immune responses is complex, and further experimental studies are needed to elucidate the mechanisms of angiotensin-mediated autoimmune renal injury.

Previous studies have shown that AT_{1A} receptor knock-out mice display marked stimulation of renin expression, increased circulating levels of angiotensin II, but with rather low-normal blood pressure. This marked activation of RAS would predict exaggerated stimulation of the residual elements of the system, including activation of other receptors of RAS. In the study by Crowley *et al.*, it was anticipated that elimination of the major angiotensin receptor AT_{1A} would be protective on murine lupus nephritis. Yet it was found that the autoimmune nephritis was accelerated. In mice, but not in humans, there is a second AT₁ receptor—AT_{1B}—whose expression was preserved in this model. As in the murine kidney, the AT_{1B} receptor is exclusively expressed by glomerular podocytes, and the AT_{1A}-deficient MRL-*Fas*^{lpr/lpr} mouse strain provides a fortuitous model wherein the renal AT₁ receptor expression is limited to the glomerulus. Thus, it could be concluded that the systemic activation of RAS in this lupus mouse line promoted more severe injury by stimulating the residual glomerular AT_{1B} receptors.

Role of RAS inhibition in immune/inflammatory renal injury

From this experimental study, one would like to extrapolate that the blockade of RAS with ACEi or ARB is very useful to protect the kidney in autoimmune injury such as in patients with lupus nephritis, or in patients with other immune complex-mediated glomerulonephritides. This has in fact been shown in previous experimental studies, and is shown again here by the fact that the ARB losartan prevented the enhanced disease process in lupus mice with AT_{1A} deficiency [1,8]. One does not need to mention the numerous studies that have documented the beneficial effects of ACEi or ARB in various glomerular diseases, including immune-mediated entities such as IgA nephritis or membranous GN, or non-immune glomerular diseases such as diabetic nephropathy.

What needs to be stated is the fact that the mechanisms by which RAS promotes immune or inflammatory glomerular damage may be far more complex than previously thought, and may not be operating solely through the AT₁ receptors. Various components of RAS, including prorenin, renin, angiotensin₁₋₇ to name a few, and the non-AT₁ receptors AT₂, Ang1-7 receptor (Mas) and others may target the immune system and renal parenchymal cells within the glomerulus in a sophisticated way, not to mention the complex interplay with other systems that include the kallikrein-bradykinin system, the NO system and the prostaglandins and thromboxanes. A dysbalance in these systems may cause different patterns of glomerular injury. Thus, one could speculate that in the future there might be more refined therapeutic strategies to influence these interacting systems. Despite decades of intense research, RAS is still always good for a surprise, and the study by Crowley *et al.* provides another elegant example of this ever increasingly complex system.

In conclusion, activated RAS has a direct negative impact on glomerular injury processes, occurring independently of its effect on blood pressure. Currently, the pharmacological blockade of RAS with ACEi or ARB, and perhaps direct renin inhibitors, appears to be the best available strategy to improve the negative impact of RAS activation on glomerular injury and loss of renal function.

Conflict of interest statement. None declared.

References

1. Crowley SD, Vasievich MP, Ruiz P *et al.* Glomerular type 1 angiotensin receptors augment kidney injury and inflammation in murine autoimmune nephritis. *J Clin Invest* 2009; 119: 943–953
2. Nataraj C, Oliverio MI, Mannon RB *et al.* Angiotensin II regulates cellular immune responses through a calcineurin-dependent pathway. *J Clin Invest* 1999; 104: 1693–1701
3. Benndorf RA, Krebs C, Hirsch-Hoffmann B *et al.* Angiotensin II type 2 receptor deficiency aggravates renal injury and reduces survival in chronic kidney disease in mice. *Kidney Int* 2009; 75: 1039–1049
4. Wolf G, Ziyadeh FN, Thaïss F *et al.* Angiotensin II stimulates expression of the chemokine RANTES in rat glomerular endothelial cells.

- Role of the angiotensin type 2 receptor. *J Clin Invest* 1997; 100: 1047–1058
5. Teplitsky V, Shoenfeld Y, Tanay A. The renin-angiotensin system in lupus: physiology, genes and practice, in animals and humans. *Lupus* 2006; 15: 319–325
6. Herlitz H, Edenö C, Mulec H *et al.* Captopril treatment of hypertension and renal failure in systemic lupus erythematosus. *Nephron* 1984; 38: 253–256
7. Pérez De Lema G, De Wit C, Cohen CD *et al.* Angiotensin inhibition reduces glomerular damage and renal chemokine expression in MRL/lpr mice. *J Pharmacol Exp Ther* 2003; 307: 275–281
8. De Albuquerque DA, Saxena V, Adams DE *et al.* An ACE inhibitor reduces Th2 cytokines and TGF- β_1 and TGF- β_2 isoforms in murine lupus nephritis. *Kidney Int* 2004; 65: 846–859

Received for publication: 11.6.09; Accepted in revised form: 11.6.09